

Ultimate Breastfeeding Prep

RESOURCE GUIDE

Helpful Links

- [AAP Statement on Breastfeeding](#)
- [Breast Massage and Hand Expression](#)
- [Getting a Deep Latch](#)
- [Breastfeeding Videos](#)
- [Multi-Language Breastfeeding Videos](#)
- [How to Store Your Breast Milk](#)
- [Breast Pump Quick Guide](#)
- [Breast Pump Cleaning Guide](#)
- [Breast Crawl](#)
- [Laid-Back Positioning](#)
- [Antenatal Expression](#)
- [Pee & Poop Chart](#)
- [Hand Expression Video](#)
- [Hand Expression Video in Spanish](#)
- [Syringe, Cup, and Finger Feeding](#)
- [SNS \(supplemental nursing system\)](#)
- [Paced Bottle Feeding](#)
- [Dietary Restrictions for Breastfeeding](#)
- [Alcohol & Breastfeeding Chart](#)
- [InfantRisk](#)
- [Lactation Amenorrhea Method \(LAM\)](#)
- [Breast Milk Volume](#)
- [My Breast Friend Nursing Pillow](#)

[**SHOP RECOMMENDED PRODUCTS**](#)



The Pumping Room

Your go-to source for all things pregnancy and breastfeeding.

[**JOIN TODAY!**](#)

Canopie Mental Health App

Canopie is your partner in maternal mental health. Aeroflow moms have free access to the app, including customized audio & video programs - plus live classes! Free access code: FEEDPREP

[**DOWNLOAD APP**](#)



Ultimate Breastfeeding Prep

RESOURCE GUIDE

Class Notes Table of Contents

- Benefits of Breastfeeding
- Golden Hour/Skin-to-Skin
- Milk Making Process
- Letdown
- Maintaining a Healthy Milk Supply
- Latch
- Positioning
- Frequency of Feeds
- Feeding Cues
- Hand Expression
- Artificial Nipples
- Diet/Alcohol and the Effect on Breastfeeding
- Partner Support
- Recommended Duration of Breastfeeding
- Lactation Amenorrhea Method
- Signs Breastfeeding is Going Well
- References



Ultimate Breastfeeding Prep

Class Notes

Benefits of Breastfeeding

- The AAP Statement on Breastfeeding includes all of the long term health benefits for mom and baby when breastfeeding is successful.

Golden Hour/Skin-to-Skin

- Helps get breastfeeding off to a great start.
- Helps with bonding and calms baby.
- Helps baby transition to the outside world and transfers good bacteria to baby.
- Reduces mom's anxiety and boosts confidence in her parenting ability.
- Stabilizes a newborn's vital signs: temperature, blood sugar, oxygen, heart rate.
- Initial assessments of baby's health can be done while mom is holding baby.
- Partners can do skin-to-skin anytime mom is not available.
- Room-in with baby after delivery to keep baby close to feed.



Milk Making Process

- **Supply and Demand**
 - The more milk that is removed, the more milk you will make.
 - Breast Milk Volume
- **Colostrum**
 - The first milk your body began making around 16 weeks of pregnancy.
 - It is small in volume, but packed with nutrition and antibodies.
 - Frequent feedings of colostrum are all baby needs just after birth.
- **Transitional Milk**
 - Most moms experience their milk “coming in” at 3-5 days after birth.
 - Breasts can feel full, heavy, warm and tender.
 - Engorgement lasts about 2-4 days.
 - Engorgement can be alleviated by frequent milk removal, cool compresses and breast massage and hand expression.
- **Mature Milk**
 - Milk is often whiter and thinner than transitional milk.
 - Milk supply continues to increase, based on demand, until about 6 weeks postpartum.

Letdown

- Also known as the milk ejection reflex.
- Milk flow is a pattern of fast and slow flows and is driven by oxytocin.
- A letdown may feel like a tingling sensation, a feeling of fullness or you notice your baby is swallowing more at the breast.



Maintaining a Healthy Milk Supply

- Nurse frequently - 8 to 12 times in 24 hours.
- Avoid artificial nipples if possible until breastfeeding is established (4-6 weeks).
- Keep baby close during the day and night.
- Prioritize feeding your baby during this very brief time that he or she is young.

Latch

- **Breastfeeding shouldn't be painful.**
 - Pain is a sign of a poor latch which can lead to nipple damage, inadequate weight gain, reduced milk supply, clogged ducts, and mastitis.
- **Keys to getting a good latch include:**
 - Baby is tummy-to-tummy with mom.
 - Align baby's nose to mom's nipple.
 - Use a "C" or "U" hold to shape and support the breast.
 - Bring baby to the breast with the head extended back.
 - Latch baby in an asymmetrical latch at the breast.
- **Signs of a good latch include:**
 - Lips are flared outward.
 - Lips are relaxed.
 - Chin touches breast.
 - Wide angle at the corner of the mouth.
 - Lots of breast and areola in the mouth.
 - No pain in the breast or nipple.



Latch (cont)

- [Getting a Deep Latch](#)
- [Breastfeeding Videos](#)
- [Multi-Language Breastfeeding Videos](#)
- If you need to pump in the early days and/or weeks postpartum while baby's latch is being addressed by an IBCLC or other provider - here are pump resources:
 - [How to Store Your Breast Milk](#)
 - [Breast Pump Quick Guide](#)
 - [Breast Pump Cleaning Guide](#)



Positioning

- Baby's first position is the **Breast Crawl**.
- Body alignment is key no matter which position you use. Be sure that baby's ears, shoulders, and hips are all in alignment.

Laid-Back - Feeding baby in a laid back or reclined position lets gravity help to keep baby close and is great for babies with latching difficulty.



Side-lying - This position helps you rest and recover while feeding your baby!



Cradle - A classic breastfeeding position where you sit upright with baby in front of you, with their head and neck laying along your forearm and their body against your stomach.



Cross Cradle - Similar to the cradle position, but one arm holds the breast while the other arm helps guide baby to the breast.



Football - An upright position where baby is placed alongside you and supported by your forearm.



Frequency of Feeds

- Newborns should breastfeed at least 8-12 times in 24 hours.
- It is NORMAL that baby does not feed on a schedule.
- Allow baby to be in control of when the feeding stops.

Feeding Cues

- Watch for mouthing, sticking out the tongue, bringing the hands to the mouth, and rooting.
- Crying is a late sign of hunger.

Hand Expression

- Hand expression can be effective at removing colostrum.
- Hand expression is a great way to express milk any time a breast pump isn't available.
- [Hand Expression Video](#)
- [Hand Expression Video in Spanish](#)

Artificial Nipples

- It is recommended to not introduce bottles and pacifiers until breastfeeding is well-established (around 4-6 weeks).
- [Syringe, cup, and finger feeding](#) can be good alternatives if supplementation is needed in the first days after birth.
- An [SNS \(supplemental nursing system\)](#) is another alternative that is useful in some cases.
- [Paced Bottle Feeding](#) is a technique that helps to protect breastfeeding and slow the flow for babies if a bottle is needed.
- Pacifiers should not be used to suppress hunger cues or delay feedings.



Diet/Alcohol and the Effect on Breastfeeding

- There are no specific **dietary restrictions for breastfeeding**.
 - Eating enough food and hydrating well is important for milk production.
- **Caffeine**
 - Up to 300mg of caffeine per day is considered safe for breastfeeding mothers and babies.
 - Caffeine may increase wakefulness and irritability in babies.
- **Alcohol**
 - It is best to allow 2 hours per drink for your body to metabolize alcohol before feeding your baby.
 - **Alcohol & Breastfeeding Chart**
- **Medications**
 - **InfantRisk** has up-to-date information regarding medications and breastfeeding.

Partner Support

- Be sure mom is eating and drinking.
- Hold baby while mom showers or naps.
- Change and bathe baby.
- Take on household duties.
- Arrange additional help (doula, housekeeper, family support).
- Ask how you can help.
- Tell her she is doing a good job.
- Anytime you are meeting your baby's needs, you are bonding.



Recommended Duration of Breastfeeding

- The AAP and WHO both recommend exclusive breastfeeding for the first 6 months before introducing solid foods, and then continuing to breastfeed for up to 2 years and beyond as desired by mother and child.

Lactation Amenorrhea Method

- Some mothers use the Lactation Amenorrhea Method (LAM) to delay fertility.
- This method must be followed exactly in order to be fully effective.
- Talk with your provider about family planning.

Signs Breastfeeding is Going Well

- Lots of wet and dirty diapers.
- Hearing and seeing soft swallows.
- Meeting growth and development milestones.
- Baby is active and alert during feedings.
- Feeding at least 8 times every 24 hours.
- Back to birth weight by 2 weeks of age.
- No breast or nipple pain.

View
Upcoming
Classes

Book an
Appointment



References

- <https://doi.org/10.1542/peds.2011-3552>
- <https://pubmed.ncbi.nlm.nih.gov/29223210/>
- <https://doi.org/10.1186/s13006-015-0035-8>
- <https://doi.org/10.1089/bfm.2013.9979>
- <https://doi.org/10.1089/bfm.2018.29110.mha>
- https://downloads.aap.org/AAP/PDF/AAP-Responsive-Feeding_Print-Fact-Sheet.pdf
- <https://www.mayoclinic.org/healthy-lifestyle/infant-and-toddler-health/expert-answers/newborn/faq-20057752>
- <https://med.stanford.edu/newborns/professional-education/breastfeeding/hand-expressing-milk.html>
- <https://doi.org/10.1089/bfm.2017.29038.ajk>
- <https://www.cdc.gov/breastfeeding-special-circumstances/hcp/diet-micronutrients/maternal-diet.html>
- <https://www.youtube.com/watch?v=OGPm5SpLxXY>
- <https://doi.org/10.1089/bfm.2015.9992>
- <https://www.cdc.gov/breastfeeding/breastfeeding-special-circumstances/vaccinations-medications-drugs/alcohol.html>
- <https://www.springerpub.com/hale-s-medications-mothers-milk-2023-9780826160638.html>
- <https://doi.org/10.1542/peds.2022-057988>
- <https://doi.org/10.1089/bfm.2014.9996>

